

RESEARCH ARTICLE

Interaction of dietary polyphenols with bovine milk proteins: Molecular structure–affinity relationship and influencing bioactivity aspects

Jianbo Xiao^{1,2}, Fangfang Mao^{1*}, Fan Yang^{1*}, Yalu Zhao¹, Chao Zhang¹ and Koichiro Yamamoto²

¹ Department of Biology, College of Life and Environment Science, Shanghai Normal University, Shanghai, P. R. China

² Department of Nutrition, Faculty of Health and Welfare, Okayama Prefectural University, Soja, Okayama, Japan

Scope: Dietary flavonoids and stilbenes are important polyphenols in foods, such as, e.g. fruits, vegetables, nuts, and tea as they are of great interest for their bioactivities, which are related to the anti-oxidative property.

Methods and results: The relationship between the structural properties of dietary polyphenols and their affinities for milk proteins (MP) was investigated. Methylation and methoxylation of flavonoids decreased (or hardly affected) the affinities for MP. Hydroxylation on the rings A and B of flavones and flavonols enhanced the interaction slightly. The hydroxylation on the ring A of flavanones significantly improved the affinities. Glycosylation of flavonoids weakened the affinities by 1–2 orders of magnitude. The hydrogenation of the C2=C3 double bond of flavonoids decreased the binding affinities by 7.24–75.86-fold. Galloylation of catechins significantly improved the binding affinities by about 100–1000-fold. Glycosylation of resveratrol decreased its affinity for MP. Esterification of gallic acid increased its binding affinity. MP significantly weakened the DPPH radical scavenging activity of polyphenols. The decreasing DPPH scavenging percentages of polyphenols increased with increasing affinities of MP–polyphenol complexes.

Conclusion: The binding affinities with MP were strongly influenced by the structural differences of dietary polyphenols. The MP–polyphenol interaction weakened with the DPPH free radical scavenging potential of polyphenols.

Received: April 25, 2011

Revised: May 4, 2011

Accepted: May 26, 2011



Keywords:

DPPH radical scavengers / Milk proteins / Polyphenols / Protein–polyphenol interaction / Structure–affinity relationship

1 Introduction

Dietary flavonoids and stilbenes are important polyphenols in foods, such as, e.g. fruits, vegetables, nuts, and tea [1–5],

as they are of great interest for their bioactivities, which are basically related to the anti-oxidative property [5–8]. The structural differences between the various classes occur in the chemistry of the ring C, as well as the number and distribution of hydroxyl groups and their substitutions on the rings A and B. These differences significantly affect their absorption, metabolism, and bio-activities in vivo. For instance, methylation of the free hydroxyl groups in the flavones dramatically increased their intestinal absorption and metabolic stability by preventing the formation of

Correspondence: Dr. Jianbo Xiao, Department of Biology, College of Life and Environment Science, Shanghai Normal University, 100 Guilin Rd, Shanghai 200234, P. R. China
E-mail: jianboxiao@yahoo.com
Fax: +86-21-64321291

Abbreviations: EC, (–)-epicatechin; EGCG, (–)-epigallocatechin gallate; GCG, gallic acid gallate; MP, milk protein

*These authors have contributed equally to this work.

glucuronic acid and sulfate conjugates [9, 10]. Walle concluded that the methylation appears to be a simple and effective way to increase the metabolic resistance and transport of flavonoids [11]. Courts and Williamson found that the C-glycoside aspalathin was methylated and glucuronidated in vivo in an intact form in human [12]. The flavonol moiety, i.e. the 2,3-double bond in conjugation with a 4-oxo group and a 3-hydroxyl group, as well as the 5,7-dihydroxylation at the A-ring has been found to be important structural features for significant anti-oxidant activity [13]. In addition to –OH moieties in the structural arrangements of flavonoids, the resonance of electrons between the rings A and B was reported to be essential for the anti-oxidant and biological activities of the compounds [14].

Milk proteins (MP) are natural vehicles, which evolved to deliver essential micronutrients (e.g. calcium and phosphate) and immune system components (e.g. immunoglobulins and lactoferrin), from mother to the newborn [15]. Briefly, cow MPs consist of caseins (2.4–2.8%), β -lactoglobulin (0.2–0.4%), α -lactalbumin (0.1–0.15), bovine serum albumin (BSA, 0.01–0.04%), immunoglobulins (0.06–0.1%), etc. [16]. Proteins from milk are natural vehicles for bioactive small molecules [15]. Many of their structural and physicochemical properties, such as excellent surface and self-assembly properties, superb gelation properties; pH-responsive gel swelling behavior, useful for programmable release; various shielding capabilities and essential for protecting sensitive payload, facilitate their functionality in binding of ions and small molecules [15, 17, 18]. Some of them (e.g. BSA, β -lactoglobulin, and γ -globulin) can bind with many kinds of endogenous and exogenous agents such as dietary polyphenols [19–24]. Few reports, however, have focused on the structure–affinity relationship of dietary polyphenols on the affinities for MP. The present work deals with the relationship between the molecular structures of dietary polyphenols and their affinities for MP. Fifty-five polyphenols (Table 1) were studied. The influence of MP–polyphenol interaction on the DPPH free radical scavenging potential of polyphenols was also investigated in detail.

2 Materials and methods

2.1 Apparatus and reagents

The fluorescence spectra were recorded on a JASCO FP-6500 fluorometer (Tokyo, Japan). The pH measurements were carried out on a Cole-Parmer PHS-3C Exact Digital pH meter (IL, USA). Biochanin A, genistein, apigenin, puerarin, catechin, (–)-epicatechin (EC), and luteolin (99.0%) were purchased from Aladin (Shanghai, China). Flavone, chrysin, and baicalein (99.5%) were obtained commercially from Wako Pure Chemical Industries (Osaka, Japan). Flavanone, 7-hydroxyflavanone, 6-hydroxyflavanone, 6-methoxyflavanone, 6-hydroxyflavone, and 6-methoxyflavone were purchased from TCI Chemical Industries (Tokyo,

Japan). Others polyphenol standards were obtained commercially from Shanghai Tauto Biotech (Shanghai, China). The working solutions of the polyphenols (1.0×10^{-3} mol/L) were prepared by dissolving each polyphenol in methanol. The pure bovine milk was obtained from Guangming (Shanghai, China). It contains (per 100 mL) 3.0 g protein, 3.2 g fat, and 4.7 g carbohydrate. The working solution of MP (1:100) was prepared by directly diluting above bovine milk with double-distilled water and stored in refrigerator prior to use. All other reagents and solvents were of analytical grade and all aqueous solutions were prepared using newly double-distilled water.

2.2 Fluorescence spectra

The fluorescence spectra were recorded upon excitation at 280 nm when MPs were titrated with polyphenols. Slit widths, scan speed, and excitation voltage were kept constant within each data set. Titrations were performed manually by using trace syringes. The fluorescence emissions of these polyphenols within the range of 300–400 nm were not observed under the excitation wavelength of 280 nm. The polyphenols were stable during the fluorescence measurements, as shown by HPLC analyses (not given here). Each fluorescence intensity determination was repeated and found to be reproducible within experimental errors.

2.3 Preparation of MP–polyphenol complexes

The working solutions of the polyphenols (1.0×10^{-3} mol/L) were diluted ten times with buffer or MP solution (1:100) to obtain tested samples. These tested samples were kept at room temperature. Samples were tested periodically for DPPH radical scavenging activity analysis.

2.4 DPPH radical scavenging activity

The anti-oxidant activities of polyphenols in the absence and presence of MP were measured on the basis of the scavenging activity of DPPH free radical. One milliliter sample was added to 1 ml of DPPH solution (0.2 mM in ethanol) as the free radical source. The decrease in the solution absorbance was measured at 517 nm after 30 min. The DPPH radical scavenging activity was calculated using the following formula:

$$\text{DPPH radical scavenging activity (\%)} = (1 - A_0/A_1) \times 100$$

where A_0 is the absorbance of the control and A_1 is the absorbance of samples.

Each experiment was repeated thrice and found to be reproducible within experimental errors.

Table 1. Chemical structures of the various polyphenols and their affinities for MP








Subclass	Name	Substitutions			Affinity		
		OH	OCH ₃	Others	lg K _a	n	
<p><i>Flavones</i></p> 	Flavone				4.81	1.05	
	7-Hydroxyflavone	7			5.09	1.05	
	6-Hydroxyflavone	6			6.02	1.16	
	6-Methoxyflavone		6		5.63	1.10	
	Chrysin	5,7			5.19	1.04	
	Baicalein	5,6,7			6.14	1.17	
	Baicalin	5,6		7-β-D-glucuronide	4.76	0.99	
	Apigenin	5,7, 4'			5.82	1.14	
	Luteolin	5,7,3', 4'			5.82	1.13	
	Hispidulin	5,7, 4'	6		5.29	1.07	
	Wogonin	5,7	8		5.13	1.07	
	Tangeretin		5,6,7,8,4'		5.76	1.12	
	Nobiletin		5,6,7,8,4',5'		5.69	1.11	
<p><i>Flavonols</i></p> 	Galangin	3,5,7			5.40	1.08	
	Kaempferide	3,5,7		4'	5.36	1.05	
	Kaempferol	3,5,7, 4'			5.57	1.06	
	Kaempferitrin	5,4'		3,7-Dirhamnoside	4.99	1.03	
	Quercetin	3,5,7,3', 4'			5.72	1.12	
	Quercitrin	5,7,3', 4'		3-O-β-D-glucoside	5.65	1.09	
	Myricetin	3,5,7,3', 4', 5'			6.04	1.17	
	Fisetin	3,7,3', 4'			5.40	1.07	
	Rutin	5,7,3', 4'		3-α-L-Rham-1,6-D-Glc	3.98	0.87	
	<p><i>Isoflavones</i></p> 	Formononetin	7	4'		4.12	0.91
		Genistein	5,7,4'			4.76	1.01
		Daidzein	7,4'			5.29	1.04
		Daidzin	4'		7-Glucoside	5.21	1.09
Genistin		5,4'		7-Glucoside	4.65	1.04	
Biochanin A		5,7	4'		4.79	0.96	
Tectorigenin		5,7,4'	6		5.47	1.11	
Puerarin		7,4'		8-C-glucose	4.21	0.90	
Sophoricoside		5,7		4'-O-glucoside	3.39	0.79	
<p><i>Flavanone</i></p> 		Naringenin	5,7, 4'			3.94	0.86
	Naringin	5,4'		7-Neohesperidose	3.76	0.83	
	Narirutin	5,4'		7-α-L-Rham-1,6-D-Glc	3.64	0.82	
	Hesperitin	5,7, 3'	4'		5.14	1.05	
	Hesperitin-7-O-rutinose	5,3'	4'	7-α-L-Rham-1,6-D-Glc	4.80	0.99	
	Dihydromyricetin	3,5,7,3', 4', 5'			5.07	1.04	
	Flavanone				3.58	0.84	
	7-Hydroxyflavanone	7			5.59	1.12	
	6-Hydroxyflavanone	6			4.87	1.08	
	6-Methoxyflavanone		6		4.77	1.06	
	Silibinin				5.28	1.07	
	Alpinetin	7	5		5.43	1.17	
	<p><i>Flavanonol</i></p> 	GCG (2,3-trans)	5,7,3',4',5'		3-Gallate	4.98	1.02
EGCG (2,3-cis)		5,7,3',4',5'		3-Gallate	4.54	0.97	
ECG (2,3-cis)		5,7,4',5'		3-Gallate	4.96	1.01	
EC (2,3-cis)		3,5,7,4',5'			3.01	0.79	
EGC (2,3-cis)		3,5,7,3',4',5'			2.05	0.65	
C (2,3-trans)		3,5,7,4',5'			2.29	0.66	
<p><i>Stilbene</i></p> 		Resveratrol	3,5,4'			4.94	1.02
		Polydatin	5,4'		3-glucoside	4.54	0.95

Table 1. Continued.

Subclass	Name	Substitutions			Affinity	
		OH	OCH ₃	Others	lg K _a	n
<i>Gallic acid</i> 	Gallic acid			R = H	5.06	1.18
	Methyl gallate			R = methyl	5.71	1.20
	Ethyl gallate			R = ethyl	5.58	1.18
	Propyl gallate			R = propyl	5.50	1.15

3 Results and discussion

3.1 The binding constants (K_a) and the number of binding sites (n)

As representative examples, the fluorescence spectra of MP after addition of daidzein and daidzin are shown in Supporting Information Fig. 1 (the fluorescence spectra of MP quenched by other polyphenols are not given here). The maximum λ_{em} of MP was hardly shifted in the presence of daidzein and daidzin. However, in these and all other cases, the fluorescence intensities of MP decreased remarkably with increasing concentration of polyphenols. The quenching degrees of polyphenols depend on their structures. For daidzein and daidzin, daidzein (A) showed stronger quenching effect on MP fluorescence than that of daidzin (B). Daidzein resulted in quenching about 48% MP fluorescence; however, daidzin only quenched about 35% MP fluorescence.

The binding constants were calculated according to the double-logarithm equation [20–25]:

$$\log_{10}(F_0 - F)/F = \log_{10}K_a + n \log_{10}[Q] \quad (1)$$

where F_0 and F represent the fluorescence intensities of MP in the absence and in the presence of polyphenols, K_a is the binding constant, n is the number of binding sites, and $[Q]$ is the concentration of polyphenols. Table 1 summarizes the calculated results according to Eq (1). The values of $\log_{10}K_a$ are proportional to the number of binding sites (n) (data not shown), which indicates that the Eq. (1) used here is suitable to study the interaction between polyphenols and MP [26, 27]. The magnitudes of apparent binding constants for MP were almost all in the range of 10^4 – 10^5 L/mol, which were similar to the recent reports for bovine γ -globulin [24]. However, these data were much smaller than the affinities of polyphenols for serum albumins from our previous reports (10^4 – 10^8 L/mol) [20–23].

3.2 Flavonoids

3.2.1 Methylation and methoxylation

As shown in Fig. 3, the methylation and methoxylation of flavonoids weakened or little affected their binding affinities




Figure 1. The methoxylation and methylation of flavonoids weakens the affinity of the polyphenols for MP.

for MP. In general, the methylation of hydroxyl group in flavonoids decreased their binding affinities for MP by 1.10–14.79 times. The affinity of daidzein for MP was found to be 14.79-times higher than that of its methylated form (formononetin) (Fig. 1). On the other side, 4'-methoxylation of galangin hardly affected the affinity for MP. This result supports that the methylation of hydroxyl group in flavonoids reduced the binding affinities for MP.

3.2.2 Hydroxylation

Table 2 shows the effects of hydroxylation of flavonoids on the affinities for MP. As seen from the data, the hydroxylation on the rings A and B of flavones and flavonols slightly enhanced the binding affinities for MP. The hydroxylation on the ring A of flavanones significantly improved the affinities for MP. However, the hydroxylation on the ring C of flavones hardly influenced the binding affinities for MP and the hydroxylation on ring A of isoflavones reduced or little affected the affinities for MP.

3.2.2.1 Hydroxylation on ring A of flavones

Hydroxylation on ring A of flavones obviously improved the affinities for MP with increasing hydroxyl group number. It

Table 2. Effects of hydroxylation of flavonoids on the affinities for MP

Class	Ring	Position	Example	Effect (times)
Flavone	A	7 H→OH	Flavone→7-OHflavone	↑ 1.91
		5 H→OH	7-OHflavone→chrysin	↑ 1.26
		6 H→OH	Chrysin→baicalein	↑ 8.91
	B	4'H→OH	Flavone→6-OHflavone	↑ 16.22
		3'H→OH	Chrysin→apigenin	↑ 4.27
		3 H→OH	Apigenin→luteolin	No effect
	C	3 H→OH	Chrysin→galangin	↑ 1.62
			Apigenin→kaempferol	↓ 1.78
			Luteolin→quercetin	↓ 1.26
Flavonol	A	5 H→OH	Fisetin→quercetin	↑ 2.09
	B	3'H→OH	Kaempferol→quercetin	↑ 1.41
		4'H→OH	Galangin→kaempferol	↑ 1.48
		5'H→OH	Quercetin→myricetin	↑ 2.09
Isoflavone	A	5 H→OH	Daidzein→genistein	↓ 3.39
			Daidzin→genistin	↓ 3.63
			Formononetin→biochanin A	Little effect
			Flavanone→6-OHflavanone	↑ 19.95
Flavanone	A	6 H→OH	Flavanone→6-OHflavanone	↑ 19.95
		7 H→OH	Flavanone→7-OHflavanone	↑ 104.71

appears that the optimal hydroxyl groups introduced to the ring A of flavones is 5,6,7-position, as the highest binding was observed with baicalein (containing three hydroxyl groups).

3.2.2.2 Hydroxylation on ring B of flavones

Hydroxylation on ring B of flavones increased or little affected the apparent binding constants (K_a) between flavones and MP. For example, the affinity of apigenin (5,7,3') for MP was found to be 4.27-times higher than that of chrysin (5,7). On the other hand, the affinity of apigenin (5,7,3') for MP was the same as that of luteolin (5,7,3',4').

3.2.2.3 Hydroxylation on ring C of flavones

The affinities of kaempferol (3,5,7,3'), galangin (3,5,7), and quercetin (3,5,7,3',4') for MP are almost the same as those of apigenin (5,7,3'), chrysin (5,7), and luteolin (5,7,3',4'), respectively. It illustrated that the hydroxylation on the ring C of flavones hardly influenced the binding affinities for MP.

3.2.2.4 Hydroxylation of rings A and B of flavonols

It was found that the hydroxylation on rings A and B of flavonols slightly improves the binding affinity for MP. The hydroxylation on ring B of flavonols obviously improved the affinities for MP with increasing hydroxyl group number. Flavonols are the most prominent flavonoids in plants. The most prominent flavonols such as quercetin and kaempferol in foods carry 5,7-dihydroxyl groups on ring A. Here, it was found that the hydroxylation on position 5 of flavonol slightly enhances the binding affinity for MP by 2.09 times.

3.2.2.5 Hydroxylation on ring A of isoflavones

The affinity of genistein (5,7,4') for MP was about 3.39-times lower than that of daidzein (7,4'). The affinity of genistin

(5,7,4') for MP was about 3.63-times lower than that of daidzein (7,4'). However, the affinity of biochanin A (5,7) for MP was similar to that of formononetin (7). These data revealed that the hydroxylation on ring A of isoflavones reduced or little affected the affinities for MP.

3.2.2.6 Hydroxylation on ring A of flavanones

Hydroxylation on ring A of flavanones significantly increased the affinities for MP. 6-OHflavanone and 7-OHflavanone showed 19.95- and 104.71-fold higher affinities than that of flavanone.

3.2.3 Comparing the affinities of flavonoid isomers with MP

In this study, we can compare two isomer groups (apigenin, galangin, baicalein, genistein and luteolin, kaempferol, fisetin). The binding constants (K_a) were determined as: baicalein > apigenin > galangin > genistein and luteolin > kaempferol > fisetin. These data illustrated that flavones showed higher binding capacity with MP than flavonols. The current results are in good agreement with data reported previously about the interaction between flavonoid isomers with BSA [28].

3.2.4 Glycosylation

The dietary flavonoids in nature occur mostly as β -glycosides. The flavonols are found mainly as the 3- and 7-O-glycoside, although the 4' position may also be glycosylated in some plants (Table 1). Most recently, we have reported the glycosylation of flavonoids lowered the affinity for BSA by 1–3 orders of magnitude depending on the conjugation




Figure 2. The glycosylation of flavonoids decreases the affinity of the polyphenols for MP. 1. K_a (quercetin)/ K_a (quercitrin), 2. K_a (daidzein)/ K_a (daidzin), 3. K_a (genistein)/ K_a (genistin), 4. K_a (naringenin)/ K_a (naringin), 5. K_a (naringenin)/ K_a (narirutin), 6. K_a (hesperitin)/ K_a (hesperitin-7-*O*-rutinose), 7. K_a (kaempferol)/ K_a (kaempferitrin), 8. K_a (daidzein)/ K_a (puerarin), 9. K_a (genistein)/ K_a (sophoricoside), 10. K_a (baicalein)/ K_a (baicalin), 11. K_a (quercetin)/ K_a (rutin).

site and the class of sugar moiety [23]. Herein, the effect of glycosylation of dietary flavonoids on the affinities for MP was investigated. The sugar moieties are in 3, 7, or 4'-positions of flavonoids. In our present study (Fig. 2), the glycosylation of flavonoids lowered the affinity for MP 1–2 order of magnitude. The affinity of quercetin for MP was about 54.95-fold higher than that of rutin, but the affinity of quercetin for MP is only 1.02 times higher than that of quercitrin. The affinities of naringenin, naringin and narirutin for MP were determined as 3.94, 3.76, and 3.64, respectively. It revealed that the monoglycosides of flavonoids showed stronger binding affinities with MP than their polyglycoside forms. The decreasing affinity for MP after glycosylation may be caused by the non-planar structure. After the hydroxyl group is substituted by a glycoside, steric hindrance may take place, which weakens the affinity for proteins [23].

3.2.5 Hydrogenation of the C2=C3 double bond

The C2=C3 double bond in conjugation with a 4-oxo group plays a very important role for the affinity for MP. As shown in Table 2, it was found that the hydrogenation of the C2=C3 double bond of flavonoids decreased the binding affinities for MP about 7.24 to 75.86 times. The affinities of apigenin and myricetin for MP were about 75.86-times and 8.51-times higher than those of naringenin and dihydromyricetin, respectively. Previously, we have investigated the effect of hydrogenation of the C2=C3 double bond in flavonoids on the affinities for BSA [29]. Hydrogenation of the C2=C3 double bond for many flavonoids decreased the binding affinity for BSA by 2–4 orders of magnitude.

Planarity of the C ring in flavonoids may be important for binding interaction with proteins, as the molecules with saturated C2–C3 bonds (flavanones and certain others) permit more twisting of the B ring with reference to the C ring. A C2=C3 double bond increases the π -conjugation of the bond linking the B and C rings, which favors near-planarity of the two rings [30]. Molecules with near-planar structure easier enter the hydrophobic pockets in proteins.

3.3 Catechins

Catechins are the major polyphenols in green tea leaves. The major catechins of green tea extract are (–)-catechin (C), EC, (–)-epigallocatechin (EGC), (–)-epicatechin gallate (ECG), (–)-epigallocatechin gallate (EGCG), and gallo catechin gallate (GCG). Here, we determined the affinities between catechins and MP by fluorescence quenching method with double-logarithm regression curve (Table 1). The binding constants ($\log_{10}K_a$) between ECG, EGCG, and GCG for MP were 4.96, 4.54, and 4.98, respectively. However, the binding constants of EC, EGC, and C for MP were within the range of 10^2 – 10^3 L/mol. It illustrates that the galloylation of catechins significantly improved the binding affinities with MP about 100–1000 times. The pyrogallol-type catechins showed lower affinities than catechol-type catechins. Moreover, the affinity of the catechin with 2,3-*trans* structure (GCG) for MP exhibited higher than that of the catechin with 2,3-*cis* structure (EGCG).

3.4 Stilbenes

Stilbenes are phytoalexins that become activated when plants are stressed and are important polyphenols with the C₆-C₂-C₆ structure [31]. The typical natural stilbenes are resveratrol and its 3-glucoside, polydatin [32]. These compounds exist in foods and are widely consumed. Resveratrol is a grape-derived polyphenol, which possesses a wide range of bioactivities including anti-oxidant, anti-inflammatory, and anti-tumor effects [33, 34]. Here, we determined the affinities between resveratrol/polydatin with MP. The binding constants ($\log_{10}K_a$) between resveratrol and polydatin for MP were 4.94 and 4.54, respectively. It illustrated that the glycosylation of resveratrol slightly reduced the affinity for MP. The affinity of resveratrol for MP was about 2.51-times higher than that of polydatin.

3.5 Gallic acid

Gallic acid is a type of phenolic acid found in gallnuts (<http://en.wikipedia.org/wiki/Gallnut>), sumac (<http://en.wikipedia.org/wiki/Sumac>), witch hazel (http://en.wikipedia.org/wiki/Witch_hazel), tea leaves (<http://en.wikipedia.org/wiki/Tea>), oak bark (<http://en.wikipedia.org/wiki/Oak>), and

other plants (<http://en.wikipedia.org/wiki/Plant>). Gallic acid seems to have anti-fungal and anti-viral properties. Gallic acid acts as an antioxidant and helps to protect our cells against oxidative damage [35]. Gallic acid was found to show cytotoxicity against cancer cells [36]. The gallates of gallic acid showed more effective inhibition of hemolysis than gallic acid [37]. The esters of gallic acid are termed as 'gallates'. The typical gallates are methyl gallate, ethyl gallate, and propyl gallate (Table 1). As seen from Table 1, the esterification of gallic acid significantly improved the affinity for MP. The affinities of gallic acid and its esters with α -amylase were determined as: methyl gallate > ethyl gallate > propyl gallate > gallic acid.

3.6 MP weakened the DPPH radical scavenging activity of polyphenols

The working solutions of the polyphenols (1.0×10^{-3} mol/L) were diluted ten times with double-distilled water or MP solution (1:100) to obtain tested samples. The immediate DPPH radical scavenging activities of the tested samples were detected to illustrate the effect of MP on the DPPH scavenging potential of polyphenols. The result showed that MP rapid weakened the DPPH radical scavenging activity of polyphenols (Fig. 3). Many reports found the similar results with the current work that proteins weakened the antioxidant capacity of polyphenols. Polyphenols–protein interaction is expected to modulate the bio-availability of polyphenols. Smith et al. found that binding of polyphenols, such as quercetin, fisetin, myricetin, and morin, to albumin reduced their prooxidant activity [38]. Arts et al. showed the antioxidant capacity of quercetin, rutin, and (+)-catechin in blood plasma was not additive [39]. Arts et al. further checked the effect of the flavonoid–protein interaction on the anti-oxidant capacity of flavonoids with the Trolox equivalent antioxidant capacity assay [40]. It looks like that




Figure 3. The DPPH radical scavenging capacity of polyphenols in the presence and absence of MP under aerobic condition.




Figure 4. The relationship between the decreased DPPH scavenging percentages (%) and $\lg K_a$ values for polyphenols.

the effect of polyphenol–protein interaction on the bio-availability of flavonoids is not equivocal and the interaction of flavonoids with proteins will weaken the antioxidant capacity of the flavonoids both in products and in vivo.

The bioactivity of polyphenols is often executed in complex biological systems where various interactions took place. Polyphenol–protein interactions are expected to modulate the bioavailability of polyphenols. The relationship of the affinities of MP–polyphenol complexes and decreasing in DPPH scavenging percentages of polyphenols in the presence of MP is shown in Fig. 4. As seen from these data, the decreasing DPPH scavenging percentages of polyphenols increased with increasing affinities of MP–polyphenol complexes. According to the free drug hypothesis [41], if a molecule is highly bound to proteins, it is difficult to disassociate this complex to form free molecule, which will weak its efficacy. Here, it looks like that the antioxidant ability of the polyphenol–MP complex does not depend on the sum of the antioxidant ability of polyphenol molecules bound but simply on the number of polyphenols unbound. The significance of current research lies in the effect of polyphenol–protein interaction on the food–drug interaction.

4 Concluding remarks

Some of the structural elements that influence the affinities of polyphenols for γ -globulin are the following: (i) Methylation and methoxylation of flavonoids decreased or little affected the affinities; (ii) Hydroxylation on the rings A and B of flavones and flavonols slightly enhanced the interaction and hydroxylation on the ring A of flavanones significantly improved the affinities; (iii) Glycosylation of polyphenols weakened the affinities; (iv) Hydrogenation of the C2 = C3 double bond of flavonoids decreased the binding affinities;

(v) Galloylation of catechins and esterification of gallic acid significantly improved the binding affinities. The MP–polyphenol interaction weakened on the DPPH free radical scavenging potential of polyphenols.

The authors are grateful for the financial support sponsored by Shanghai Rising-Star Program (11QA1404700), Natural Science Foundation of Shanghai (10ZR1421700), Leading Academic Discipline Project of Shanghai Municipal Education Commission (J50401), Innovation Program of Shanghai Municipal Education Commission (10YZ68), “Chen Guang” project supported by Shanghai Municipal Education Commission and Shanghai Education Development Foundation (09CG46), Program of Shanghai Normal University (SK201006).

The authors have declared no conflict of interest.

5 References

- [1] Cohen, S. D., Kennedy, J. A., Plant metabolism and the environment: Implications for managing phenolics. *Crit. Rev. Food Sci.* 2010, 50, 620–643.
- [2] Hagl, S., Deusser, H., Soyalan, B., Janzowski, C. et al., Colonic availability of polyphenols and D-(–)-quinic acid after apple smoothie consumption. *Mol. Nutr. Food Res.* 2011, 55, 368–377.
- [3] Côté, J., Caillet, S., Doyon, G., Sylvain, J. F., Lacroix, M., Analyzing cranberry bioactive compounds. *Crit. Rev. Food Sci.* 2010, 50, 872–888.
- [4] Johnson, J., de Mejia, E. G., Dietary factors and pancreatic cancer: the role of food bioactive compounds. *Mol. Nutr. Food Res.* 2011, 55, 58–73.
- [5] Côté, J., Caillet, S., Doyon, G., Sylvain, J. F., Lacroix, M., Bioactive compounds in cranberries and their biological properties. *Crit. Rev. Food Sci.* 2010, 50, 666–679.
- [6] Brand, W., Padilla, B., van Bladeren, P. J., Williamson, G., Rietjens, I. M. C. M., The effect of co-administered flavonoids on the metabolism of hesperetin and the disposition of its metabolites in Caco-2 cell monolayers. *Mol. Nutr. Food Res.* 2010, 54, 851–860.
- [7] Blade, C., Arola, L., Salvado, M. J., Hypolipidemic effects of proanthocyanidins and their underlying biochemical and molecular mechanisms. *Mol. Nutr. Food Res.* 2010, 54, 37–59.
- [8] Morabito, G., Trombetta, D., Brajendra, K. S., Ashok, K. P. et al., Antioxidant properties of 4-methylcoumarins in vitro cell-free systems. *Biochimie* 2010, 92, 1101–1107.
- [9] Wen, X., Walle, T., Methylated flavonoids have greatly improved intestinal absorption and metabolic stability. *Drug Metab. Dispos.* 2006, 34, 1786–1792.
- [10] Walle, T., Methylation of dietary flavones increases their metabolic stability and chemopreventive effects. *Int. J. Mol. Sci.* 2009, 10, 5002–5019.
- [11] Walle, T., Methylation of dietary flavones greatly improves their hepatic metabolic stability and intestinal absorption. *Mol. Pharmaceut.* 2007, 4, 826–832.
- [12] Courts, F. L., Williamson, G., The C-glycosyl flavonoid, aspalathin, is absorbed, methylated and glucuronidated intact in humans. *Mol. Nutr. Food Res.* 2009, 53, 1104–1111.
- [13] Woodman, O. L., Meeker, W. F., Boujaoude, M., Vasorelaxant and antioxidant activity of flavonols and flavones: structure activity relationships. *J. Cardio Vasc. Pharm.* 2005, 46, 302–309.
- [14] Heijnen, C. G. M., Haenen, G. R. M. M., van Acker, F. A. A., van der Vijgh, W. J. F., Bast, A., Flavonoids as peroxynitrite scavengers: the role of the hydroxyl groups. *Toxicol. In Vitro* 2001, 15, 3–6.
- [15] Livney, Y. D., Milk proteins as vehicles for bioactives. *Curr. Opin. Colloid Interface Sci.* 2010, 15, 73–83.
- [16] Fox, P. F., McSweeney, P. L. H., *Advanced Dairy Chemistry, Proteins Part A, Vol. 1*, 3rd Edn, Kluwer Academic/Plenum Publishers, New York 2003.
- [17] Singh, H., The milk fat globule membrane – a biophysical system for food applications. *Curr. Opin. Colloid Interface Sci.* 2006, 11, 154–163.
- [18] Singh, H., Ye, A., in: Thompson, A., Boland, M. J., Singh, H. (Eds.), *Milk Proteins, from Expression to Food*, 1st Edn, Elsevier, Academic Press, Amsterdam 2009, 321–345.
- [19] Bae, M. J., Ishii, T., Minoda, K., Kawada, Y. et al., Albumin stabilizes (–)-epigallocatechin gallate in human serum: binding capacity and antioxidant property. *Mol. Nutr. Food Res.* 2009, 53, 709–715.
- [20] Xiao, J. B., Cao, H., Wang, Y. F., Yamamoto, K., Wei, X. L., Structure-affinity relationship of flavones on binding to serum albumins: effect of hydroxyl groups on ring A. *Mol. Nutr. Food Res.* 2010, 54, S253–S260.
- [21] Xiao, J. B., Chen, T. T., Cao, H., Chen, L., Yang, F., Molecular property-affinity relationship of flavanoids and flavonoids for human serum albumin in vitro. *Mol. Nutr. Food Res.* 2011, 55, 310–317.
- [22] Xiao, J. B., Suzuki, M., Jiang, X. Y., Chen, X. Q. et al., Influence of B-ring hydroxylation on interactions of flavonols with bovine serum albumin. *J. Agric. Food Chem.* 2008, 56, 2350–2356.
- [23] Xiao, J. B., Cao, H., Wang, Y. F., Zhao, J. Y., Wei, X. L., Glycosylation of dietary flavonoids decreases the affinities for plasma protein. *J. Agric. Food Chem.* 2009, 57, 6642–6648.
- [24] Xiao, J. B., Kai, G. Y., Yang, F., Liu, C. X. et al., Molecular structure-affinity relationship of natural polyphenols for bovine γ -globulin. *Mol. Nutr. Food Res.* 2011, 55, S86–S92.
- [25] Xiao, J. B., Cao, H., Chen, T. T., Yang, F. et al., Molecular property-binding affinity relationship of flavonoids for common rat plasma proteins in vitro. *Biochimie* 2011, 93, 134–140.
- [26] Berezhkovsky, L. M., On the calculation of the concentration dependence of drug binding to plasma proteins with multiple binding sites of different affinities: determination of the possible variation of the unbound drug fraction and calculation of the number of binding sites of the protein. *J. Pharm. Sci.* 2007, 96, 249–257.

- [27] Xiao, J. B., Chen, X. Q., Jiang, X. Y., Hilczer, M., Tachiya, M., Probing the interaction of trans-resveratrol with bovine serum albumin: a fluorescence quenching study with Tachiya model. *J. Fluoresc.* 2008, 18, 671–678.
- [28] Cao, H., Liu, Q., Shi, J., Xiao, J. B., Xu, M., Comparing the affinities of flavonoid isomers with protein by fluorescence spectroscopy. *Anal. Lett.* 2008, 41, 521–532.
- [29] Liu, X., Chen, X. Q., Xiao, J. B., Zhao, J. Y. et al., Effect of hydrogenation on ring C of flavonols on their affinity for bovine serum albumin. *J. Solution Chem.* 2010, 39, 533–542.
- [30] Edenharder, R., von Petersdorff, I., Rauscher, R., Anti-mutagenic effects of flavonoids, chalcones and structurally related compounds on the activity of 2-amino-3-methylimidazo[4,5-f]quinoline (IQ) and other heterocyclic amine mutagens from cooked food. *Mutat. Res.* 1993, 287, 261–274.
- [31] Wang, T. T. Y., Schoene, N. W., Kim, Y. S., Mizuno, C. S., Rimando, A. M., Differential effects of resveratrol and its naturally occurring methylether analogs on cell cycle and apoptosis in human androgen-responsive LNCaP cancer cells. *Mol. Nutr. Food Res.* 2010, 54, 335–344.
- [32] Zhou, S. Y., Yang, R. T., Teng, Z. H., Zhang, B. L. et al., Dose-dependent absorption and metabolism of trans-polydatin in rats. *J. Agric. Food Chem.* 2009, 57, 4572–4579.
- [33] Weng, Y. L., Liao, H. F., Li, A. F. Y., Chang, J. C., Chiou, R. Y. Y., Oral administration of resveratrol in suppression of pulmonary metastasis of BALB/c mice challenged with CT26 colorectal adenocarcinoma cells. *Mol. Nutr. Food Res.* 2010, 54, 259–267.
- [34] Cottart, C. H., Nivet-Antoine, V., Laguillier-Morizot, C., Beaudeau, J. L., Resveratrol bioavailability and toxicity in humans. *Mol. Nutr. Food Res.* 2010, 54, 7–16.
- [35] Chuang, C. Y., Liu, H. C., Wu, L. C., Chen, C. Y. et al., Gallic acid induces apoptosis of lung fibroblasts via a reactive oxygen species-dependent ataxia telangiectasia mutated-p53 activation pathway. *J. Agric. Food Chem.* 2010, 58, 2943–2951.
- [36] Forester, S. C., Waterhouse, A. L., Gut metabolites of anthocyanins, gallic Acid, 3-O-methylgallic acid, and 2,4,6-trihydroxybenzaldehyde, inhibit cell proliferation of Caco-2 Cells. *J. Agric. Food Chem.* 2010, 58, 5320–5327.
- [37] Ximenes, V. F., Lopes, M. G., Petronio, M. S., Regasini, L. O. et al., Inhibitory effect of gallic acid and its esters on 2,2'-azobis(2-amidinopropane)hydrochloride (AAPH)-induced hemolysis and depletion of intracellular glutathione in erythrocytes. *J. Agric. Food Chem.* 2010, 58, 5355–5362.
- [38] Smith, C., Halliwell, B., Aruoma, O., Protection by albumin against the pro-oxidant actions of phenolic dietary components. *Food Chem. Toxicol.* 1992, 30, 483–489.
- [39] Arts, M. J. T. J., Haenen, G. R. M. M., Voss, H. P., Bast, A., Masking of antioxidant capacity by the interaction of flavonoids with protein. *Food Chem. Toxicol.* 2001, 39, 787–791.
- [40] Arts, M. J. T. J., Haenen, G. R. M. M., Wilms, L. C., Beetstra, S. A. J. N. et al., Interactions between flavonoids and proteins: Effect on the total antioxidant capacity. *J. Agric. Food Chem.* 2002, 50, 1184–1187.
- [41] Buxton, I., in: Goodman & Gilman's The Pharmacological Basis of Therapeutics. Brunton L. (Ed.), McGraw-Hill, Chicago 2005, pp. 1–39.