

## Coffee consumption but not green tea consumption is associated with adiponectin levels in Japanese males

T. Imatoh · S. Tanihara · M. Miyazaki ·  
Y. Momose · Y. Uryu · H. Une

Received: 11 May 2010 / Accepted: 29 September 2010 / Published online: 16 October 2010  
© Springer-Verlag 2010

### Abstract

**Purpose** Coffee is among the most widely consumed beverages in the world. Numerous epidemiological studies have reported a significant inverse association between coffee consumption and risk of type 2 diabetes mellitus, but the underlying mechanisms are still not fully understood. Therefore, we conducted an epidemiological study to clarify the relationship between coffee consumption and adiponectin levels in Japanese males. We also evaluated whether green tea consumption affected adiponectin levels. **Methods** We carried out a cross-sectional study. The subjects were 665 male employees in Japan. Coffee consumption was assessed, using a self-administered questionnaire, as the number of times per week and cups per day respondents drank, and subjects were grouped into four levels (non, 1–5 times/week, 1–2 cups/day and  $\geq 3$  cups/day). **Results** The means of adiponectin levels were positively associated with coffee consumption. A dose–response relationship was found between coffee consumption and circulating adiponectin levels. The relationship remained significant after adjustment for potential confounding factors ( $P$  for trend  $<0.05$ ). However, green tea consumption was not significantly associated with adiponectin levels ( $P$  for trend = 0.90).

**Conclusions** We not only revealed that habitual coffee consumption is associated with higher adiponectin levels in Japanese males but also found a dose-dependent association between coffee consumption and adiponectin levels. Therefore, our study suggested that coffee components might play an important role in the elevation of adiponectin level.

**Keywords** Adiponectin · Epidemiological study · Coffee · Green tea

### Introduction

Coffee is among the most widely consumed beverages in the world [1]. It is believed to have beneficial effects in the prevention and treatment of many diseases. The protective effect of coffee against many diseases such as various types of cancer [2, 3], cardiovascular diseases [4] and type 2 diabetes mellitus [5–8] has been investigated.

Adiponectin is an adipocyte-specific secretory protein [9]. Circulating adiponectin levels are lower in the obese and in type 2 diabetes subjects than in healthy controls. Many studies have demonstrated that adiponectin plays an important role in the regulation of insulin function and the development of type 2 diabetes mellitus [10]. To date, a few studies have reported a finding of this sort: Williams et al. [11] found that women drinking four or more cups of coffee per day had significantly higher adiponectin levels than those who did not drink coffee regularly. Kempf et al. [12] indicated a significant increase in adiponectin levels in response to increased coffee consumption in an intervention trial.

Recent studies have demonstrated that coffee consumption can affect blood pressure, cholesterol levels and

T. Imatoh (✉) · S. Tanihara · Y. Momose · Y. Uryu · H. Une  
Department of Hygiene and Preventive Medicine,  
School of Medicine, Fukuoka University, 7-45-1,  
Nanakuma, Jonan-ku,  
Fukuoka City, Fukuoka 814-0180, Japan  
e-mail: imatoh@cis.fukuoka-u.ac.jp

M. Miyazaki  
Saitama City Institute of Health Science and Research,  
7-5-12 Suzuya Chuou-ku, Saitama City,  
Saitama 338-0013, Japan

liver transaminase levels. Therefore, it might be expected that coffee consumption would affect adiponectin levels. It is possible to hypothesize that coffee may reduce the risk for certain chronic diseases, including diabetes mellitus, through its effects on glucose metabolism and lipid metabolism, and its protective action on the liver.

This study was conducted for the purpose of clarifying the associations of coffee consumption with adiponectin levels in Japanese males. We evaluated whether coffee consumption was associated with adiponectin, metabolic parameters including parameters of liver damage, lipoprotein and white blood cells. Moreover, the current study on adiponectin levels in Japanese males examined not only associations with coffee consumption but also associations with consumption of green tea.

## Subjects and methods

### Study subjects

Study subjects were male employees registered with a particular health insurance society in Fukuoka Prefecture, Japan. All employees were full-time, white-collar, daytime Japanese workers. Each subject received an annual health check-up in 2000. The study protocol was approved by the Ethics Committee of Fukuoka University. Consent was obtained from 671 people to use the data obtained during their annual health check-ups and to donate residual blood samples used for the examination. Individuals who had missing data for any of the exposure parameters were excluded ( $n = 6$ ). Finally, 665 subjects were included in the present analysis.

### Variables

Height and weight were measured to calculate body mass index ( $BMI = \text{weight in kilograms divided by the square of height in meters}$ ). The systolic blood pressure (SBP) and diastolic blood pressure (DBP) were measured once using a standard mercury sphygmomanometer with the cuff on the right arm and the subjects in a sitting position. Smoking status, alcohol drinking status, physical activity, and coffee and green tea consumption were assessed through a self-administered questionnaire. Coffee or green tea consumptions were categorized into four levels, respectively (coffee: non, 1–5 times/week, 1–2 cups/day and  $\geq 3$  cups/day; green tea: non, 1–5 times/week, 1–3 cups/day and  $\geq 4$  cups/day). The standard serving size was assigned on the questionnaire as 1 cup. Participants were classified as never, ex-, and current smokers. Alcohol-drinking status was classified into four categories: never, ex-, 1–6 times/week and regular

alcohol drinkers. We defined physically activity as over 30 min of leisure time physical activity. Frequency of physical activity was divided two categories by the number of times (less than once a week, once a week or more). Hypertension was defined as a SBP of  $\geq 140$  mmHg and/or a DBP of  $\geq 90$  mmHg and/or taking antihypertensive medication. Hypercholesterolemia was defined as total cholesterol (TC) of  $\geq 240$  mg/dL and/or taking anti-hyperlipidemic medication. Type 2 diabetes mellitus was defined as taking anti-diabetic medication.

Adiponectin levels were determined in 2006 in spare serum samples that had been stored at  $-80^\circ\text{C}$ . All of the subjects agreed to have their serum adiponectin levels measured. Adiponectin levels were determined by enzyme-linked immunosorbent assay (Otsuka Pharmaceutical Co., Ltd, Japan). Serum total cholesterol and LDL cholesterol (LDL-C) were measured enzymatically using commercial enzyme kits (Wako, Osaka, Japan and Daiichi Kagaku, Tokyo, Japan, respectively). HDL cholesterol (HDL-C) was measured using the direct method. Triglyceride (TG) was measured using the glycerol removal method. White blood cell counts were measured by flow cytometry. Aspartate aminotransferase (AST), alanine aminotransferase (ALT) and  $\gamma$ -glutamyl transpeptidase (GGT) were measured using the Japan Society of Clinical Chemistry (JSCC) consensus method. The basic characteristics of the subjects are shown in Table 1.

### Statistical analysis

AST, ALT, GGT, TG and serum adiponectin levels were converted into logarithms to approximately normalize the distribution and used in one-way analysis of variance (ANOVA) and analysis of covariance (ANCOVA). Comparison of continuous variables including adiponectin levels against coffee-drinking status was conducted using ANOVA. Post hoc comparison was conducted using Tukey's method. ANCOVA was used to test differences and to estimate adjusted means across categories of coffee-drinking status or green tea drinking status after adjustment for covariates. These estimated means were adjusted for age (continuous), BMI (continuous), smoking status (never, ex-, current), physical activity (less than once a week, once a week or more) and taking anti-hypertensive medication (yes or no). The geometric means with adjustment for covariates are shown. Trends were tested for using the contrast method. Categorical variables were compared using the chi-square test. A  $P$  value of less than 0.05 was considered statistically significant. All statistical analyses were conducted using SAS for WINDOWS version 9.1 (SAS Institute, Cary, NC).

**Table 1** Basic characteristics of participants

| Variables                                       | <i>n</i> = 665 |
|-------------------------------------------------|----------------|
| Age (years), mean (SD)                          | 48.3 (5.3)     |
| Body mass index (kg/m <sup>2</sup> ), mean (SD) | 23.4 (2.9)     |
| Systolic blood pressure (mmHg), mean (SD)       | 125.0 (14.6)   |
| Diastolic blood pressure (mmHg), mean (SD)      | 78.0 (10.7)    |
| Smoking status, <i>n</i> (%)                    |                |
| Non-smoker                                      | 196 (43.1)     |
| Ex-smoker                                       | 57 (11.5)      |
| Current smoker                                  | 412 (45.4)     |
| Drinking status, <i>n</i> (%)                   |                |
| Alcohol                                         |                |
| Non-drinker                                     | 112 (16.8)     |
| Ex-drinker                                      | 10 (1.5)       |
| 1–6 times/week                                  | 158 (23.8)     |
| Regular drinker                                 | 385 (57.9)     |
| Coffee                                          |                |
| Non-drinker                                     | 130 (19.5)     |
| 1–5 times/week drinker                          | 181 (27.2)     |
| 1–2 cups/day drinker                            | 220 (33.1)     |
| ≥3 cups/day drinker                             | 134 (20.2)     |
| Green tea                                       |                |
| Non-drinker                                     | 113 (17.0)     |
| 1–5 times/week drinker                          | 113 (17.0)     |
| 1–3 cups/day drinker                            | 267 (40.2)     |
| ≥4 cups/day drinker                             | 172 (25.9)     |
| Physical activity, <i>n</i> (%)                 |                |
| Less than once a week                           | 522 (78.5)     |
| Once a week or more                             | 143 (21.5)     |
| Hypertension, <i>n</i> (%)                      | 112 (16.8)     |
| Hypercholesterolemia, <i>n</i> (%)              | 185 (27.8)     |
| Diabetes mellitus, <i>n</i> (%)                 | 27 (4.1)       |

## Results

The basic characteristics of subjects in relation to their coffee consumption are shown in Table 2. Subjects who consumed more coffee tended to be younger. Increasing coffee consumption was associated with lower SBP and DBP. Coffee-drinking was associated with smoking and drinking status. Current smokers and non-alcohol drinkers were more likely to consume coffee frequently. There were no significant differences between coffee consumption and BMI or frequency of physical activity.

Biological characteristics by consumption of coffee are shown in Table 3. Higher coffee consumption was significantly associated with lower levels of AST, ALT and GGT. As coffee consumption increased, white blood cell counts were higher. There were no associations between coffee consumption and TC, LDL-C or HDL-C levels. The

respective means of adiponectin in those drinking coffee non-, 1–5 times/week, 1–2 cups/day and ≥3 cups/day were, respectively, 5.95, 6.51, 7.05 and 6.89 µg/mL. The difference was significant ( $P < 0.01$ , one-way ANOVA).

Next, we performed ANCOVA to estimate adjusted means and to test the differences (Table 4). The logarithm-transformed means of adiponectin level rose as coffee consumption increased. The means were positively associated with coffee consumption even after confounding factors, such as age, BMI, smoking status, anti-hypertensive medication and physical activity, were adjusted for ( $P < 0.01$ ,  $P$  for trend  $< 0.05$ ). Though a significant dose–response relationship was observed between coffee consumption and adiponectin levels, there were no significant differences in adiponectin levels between 1 and 2 cups/days and ≥3 cups/day ( $P = 0.58$ ). On the other hand, drinking green tea was not associated with higher adiponectin levels ( $P = 0.89$ ,  $P$  for trend = 0.90).

## Discussion

This study was conducted to examine the association between coffee consumption and circulating adiponectin level. The main findings of the present study are that the adiponectin levels of those drinking 1 or more cups of coffee per day were significantly higher than those of subjects who drank coffee occasionally or never. Moreover, a significant dose–response relationship was observed.

Some previous studies demonstrated a relationship between coffee consumption and adiponectin level. Williams et al. [11] reported that subjects with and without diabetes who drank four or more cups of coffee per day had significantly higher adiponectin levels than those who did not drink coffee regularly. Moreover, similar findings were observed for caffeine intake. The study included women only. Kempf et al. [12] reported that coffee consumption led to increased adiponectin levels in an interventional study. Although adiponectin strongly correlated with BMI, the interventional study did not consider weight change. Though their studies had several limitations, our findings are consistent with theirs and suggest that coffee has a beneficial effect on blood adiponectin levels in Japanese men.

Moreover, we investigated whether coffee consumption was related to metabolic parameters. Our results indicated that higher coffee consumption was associated with lower levels of parameters of liver damage, including AST, ALT and GGT. Several studies reported that coffee consumption inhibited elevation of liver transaminases [13] and was associated with a lower risk of liver cirrhosis [14]. Although our study did not observe the association between

**Table 2** Basic characteristics by consumption of coffee

| Variables                                       | Non<br>(n = 130) | 1–5 times/week<br>(n = 181) | 1–2 cups/day<br>(n = 220) | ≥3 cups/day<br>(n = 134) | P value          |
|-------------------------------------------------|------------------|-----------------------------|---------------------------|--------------------------|------------------|
| Age (year), mean (SD)                           | 49.9 (5.0)       | 48.0 (5.4)                  | 47.7 (5.3)                | 48.2 (5.1)               | <b>&lt;0.001</b> |
| Body mass index (kg/m <sup>2</sup> ), mean (SD) | 23.7 (2.9)       | 23.8 (3.1)                  | 23.1 (2.8)                | 23.2 (2.9)               | 0.06             |
| Systolic blood pressure (mmHg), mean (SD)       | 129.5 (13.6)     | 124.6 (13.7)                | 123.1 (15.2)              | 124.6 (14.8)             | <b>&lt;0.001</b> |
| Diastolic blood pressure (mmHg), mean (SD)      | 80.7 (9.7)       | 77.7 (10.6)                 | 76.7 (10.7)               | 77.9 (11.1)              | <b>&lt;0.01</b>  |
| Smoking status, n (%)                           |                  |                             |                           |                          | <b>&lt;0.001</b> |
| Non-smoker                                      | 56 (43.1)        | 69 (38.1)                   | 54 (24.6)                 | 17 (12.7)                |                  |
| Ex-smoker                                       | 15 (11.5)        | 14 (7.7)                    | 18 (8.2)                  | 10 (7.5)                 |                  |
| Current smoker                                  | 59 (45.4)        | 98 (54.1)                   | 148 (67.3)                | 107 (79.9)               |                  |
| Alcohol drinking status, n (%)                  |                  |                             |                           |                          | <b>&lt;0.05</b>  |
| Non-drinker                                     | 13 (10.0)        | 24 (13.3)                   | 42 (19.1)                 | 33 (24.6)                |                  |
| Ex-drinker                                      | 2 (1.5)          | 2 (1.1)                     | 4 (1.8)                   | 2 (1.5)                  |                  |
| 1–6 times/week                                  | 26 (20.0)        | 52 (28.7)                   | 48 (21.8)                 | 32 (23.9)                |                  |
| Regular drinker                                 | 89 (68.5)        | 103 (56.9)                  | 126 (57.3)                | 67 (50.0)                |                  |
| Physical activity, n (%)                        |                  |                             |                           |                          | 0.72             |
| Less than once a week                           | 103 (79.2)       | 141 (77.9)                  | 177 (80.5)                | 101 (75.4)               |                  |
| Hypertension, n (%)                             | 38 (29.2)        | 22 (12.2)                   | 33 (15.0)                 | 19 (14.2)                | <b>&lt;0.001</b> |
| Hypercholesterolemia, n (%)                     | 40 (30.8)        | 51 (28.2)                   | 59 (26.8)                 | 35 (26.1)                | 0.83             |
| Diabetes mellitus, n (%)                        | 7 (5.4)          | 3 (1.7)                     | 12 (5.5)                  | 5 (3.7)                  | 0.22             |

P value by one-way analysis of variance for continuous variables and  $\chi^2$  tests for categorical variables

**Table 3** Biological characteristics by consumption of coffee

| Variables                          | Non<br>(n = 130) | 1–5 times/week<br>(n = 181) | 1–2 cups/day<br>(n = 220) | ≥3 cups/day<br>(n = 134) | P value          |
|------------------------------------|------------------|-----------------------------|---------------------------|--------------------------|------------------|
| AST (IU/L), mean (SD)              | 34.6 (25.8)      | 29.1 (14.8)                 | 28.5 (16.0)               | 29.1 (17.6)              | <b>&lt;0.01</b>  |
| ALT (IU/L), mean (SD)              | 98.0 (97.1)      | 77.8 (82.9)                 | 62.0 (68.4)               | 60.4 (75.7)              | <b>&lt;0.01</b>  |
| GGT (IU/L), mean (SD)              | 37.4 (22.4)      | 34.2 (37.8)                 | 32.1 (24.7)               | 32.3 (36.6)              | <b>&lt;0.001</b> |
| Cholesterol level (mg/dL)          |                  |                             |                           |                          |                  |
| TC, mean (SD)                      | 199.9 (34.4)     | 202.7 (33.3)                | 200.1 (33.5)              | 202.9 (35.0)             | 0.78             |
| LDL-C, mean (SD)                   | 103.2 (30.5)     | 109.2 (29.0)                | 110.3 (26.8)              | 110.8 (28.7)             | 0.10             |
| HDL-C, mean (SD)                   | 61.9 (14.7)      | 59.0 (15.2)                 | 60.3 (14.5)               | 58.3 (14.0)              | 0.17             |
| Triglyceride (mg/dL), mean (SD)    | 195.8 (169.0)    | 203.0 (210.8)               | 156.3 (113.8)             | 194.5 (312.3)            | <b>&lt;0.01</b>  |
| White blood cell count (/μL), mean | 6420.8 (1689.5)  | 6577.4 (1749.2)             | 6818.6 (1734.0)           | 7164.2 (1884.4)          | <b>&lt;0.01</b>  |
| Adiponectin level (μg/mL)          |                  |                             |                           |                          | <b>&lt;0.01</b>  |
| Mean (SD)                          | 6.0 (2.6)        | 6.5 (3.0)                   | 7.1 (3.2)                 | 6.9 (3.3)                |                  |
| Median (IQR)                       | 5.3 (4.1–7.3)    | 6.0 (4.3–8.0)               | 6.5 (4.6–8.7)             | 6.2 (4.5–8.6)            |                  |

ALT Alanine aminotransferase, AST aspartate aminotransferase, GGT  $\gamma$ -glutamyl transpeptidase, TC total cholesterol, LDL-C LDL cholesterol, HDL-C HDL cholesterol, IQR inter quartile range

P values refer to overall differences across groups as derived from one-way analysis of variance

coffee consumption and cholesterol levels, recent studies showed that coffee consumption increased HDL-C levels [12, 15].

We also examined whether green tea consumption was associated with higher adiponectin levels. Green tea is a beverage that is commonly consumed in Japan. Though

some animal studies [16, 17] have demonstrated that green tea extract can increase the levels of adiponectin, in human studies [18–20], the effects of green tea on adiponectin levels are not clear. In our study, no association between green tea consumption and adiponectin levels was found. Ryu et al. [20] reported that serum adiponectin levels were

**Table 4** Crude and adjusted means of adiponectin levels by consumption of coffee

| <i>n</i> (%)                |            | Adiponectin level (µg/mL) |                 |                 |
|-----------------------------|------------|---------------------------|-----------------|-----------------|
|                             |            | Crude                     | Model 1         | Model 2         |
| Coffee consumption          |            |                           |                 |                 |
| Non                         | 130 (19.6) | 5.47                      | 5.49            | 5.48            |
| 1–5 times/week              | 181 (27.2) | 5.91                      | 6.03            | 6.03            |
| 1–2 cup/day                 | 220 (33.1) | 6.43                      | 6.33            | 6.32            |
| ≥3 cups/day                 | 134 (20.2) | 6.19                      | 6.16            | 6.17            |
| <i>P</i> value <sup>a</sup> |            | <b>&lt;0.01</b>           | <b>&lt;0.01</b> | <b>&lt;0.01</b> |
| <i>P</i> for trend          |            | <b>&lt;0.01</b>           | <b>&lt;0.05</b> | <b>&lt;0.05</b> |
| Green tea consumption       |            |                           |                 |                 |
| Non                         | 113 (17.0) | 6.09                      | 6.16            | 6.15            |
| 1–5 times/week              | 113 (17.0) | 5.82                      | 5.91            | 5.90            |
| 1–3 cup/day                 | 267 (40.2) | 5.99                      | 6.04            | 6.04            |
| ≥4 cups/day                 | 172 (25.9) | 6.23                      | 6.05            | 6.06            |
| <i>P</i> value <sup>a</sup> |            | 0.61                      | 0.89            | 0.89            |
| <i>P</i> for trend          |            | 0.55                      | 0.82            | 0.90            |

Geometric means are shown

Model 1: Age (continuous), Body mass index (continuous), Smoking status (never, ex-, current)

Model 2: Age (continuous), Body mass index (continuous), Smoking status (never, ex-, current), Anti-hypertensive medication (yes or no), Physical activity (less than once a week, once a week or more)

<sup>a</sup> *P* value by one-way ANOVA or ANCOVA

unchanged after long-term green tea consumption [20]. Our data support their findings.

Numerous recent epidemiological studies indicate that regular coffee intake may have beneficial effects including decreased risk of type 2 diabetes mellitus [5–7, 21–23]. Several plausible mechanisms for the beneficial effect of coffee have been examined. The possibility that coffee lowers the levels of inflammatory markers such as C-reactive protein has been considered [24, 25]. It has also been suggested that coffee has antioxidant activities. It was reported that caffeine is the most plausible functional component in regular coffee to cause the increase in serum adiponectin levels in women. Since coffee contains about twice as much caffeine as green tea [26], the observed difference of adiponectin levels between coffee and green tea may depend on the caffeine content. However, it was reported that decaffeinated coffee consumption was associated with a lower risk of type 2 diabetes [27]. Apart from caffeine, coffee contains hundreds of other functional components, including phenolic compounds (chlorogenic acid and quinides), minerals (magnesium, potassium, manganese and chromium), vitamins (niacin) and fibers [28, 29]. Hence, the other components of coffee except caffeine may have beneficial effects. The exact underlying mechanisms are not yet completely clear.

There are several limitations to our study. First, we have no data about other lifestyle factors such as diet. Heavy coffee consumption tends to be strongly associated with cigarette smoking and, in many populations, with a generally less health-conscious lifestyle and lower socioeconomic status [30]. Though it is important to take into account the possible confounding role of other lifestyle factors, our study subjects had relatively similar lifestyle and socioeconomic factors. Therefore, we considered that the effects of residual confounding are relatively small. Second, we have no data on types of coffee, or whether sugar, milk or cream was added. Because the amount of sugar and milk added to coffee is probably insignificant compared to the consumption of other food, it seems that these are unlikely to have affected the metabolism. Inverse association with risk of type 2 diabetes mellitus was observed for coffee with or without milk/cream [31]. Third, because ours was a cross-sectional study, we did not deal with any temporal relationship between higher serum adiponectin levels and coffee consumption.

In conclusion, coffee consumption was found to be related to adiponectin levels in Japanese males. Adiponectin levels increased with higher coffee consumption in a graded manner. On the other hand, the benefits of green tea consumption on adiponectin levels were not confirmed in our study. The consumption of coffee, but not of green tea, appears to have benefits to adiponectin levels. Further studies are necessary to confirm our results in various populations and to determine whether the benefits of coffee consumption to circulating adiponectin level are causal.

**Acknowledgments** This work was supported by a Grant-in-Aid for Scientific Research from the Japan Society for the Promotion of Science [20590619], [21790564].

**Conflict of interest** The authors declare no conflict of interest.

## References

- Schaefer B (2004) Coffee consumption and type 2 diabetes mellitus. *Ann Intern Med* 141:323–324
- Bravi F, Bosetti C, Tavani A, Bagnardi V, Gallus S, Negri E, Franceschi S, La Vecchia C (2007) Coffee drinking and hepatocellular carcinoma risk: a meta-analysis. *Hepatology* 46:430–435
- Lee JE, Hunter DJ, Spiegelman D et al (2007) Intakes of coffee, tea, milk, soda and juice and renal cell cancer in a pooled analysis of 13 prospective studies. *Int J Cancer* 121:2246–2253
- Greenberg JA, Dunbar CC, Schnoll R, Kokolis R, Kokolis S, Kassotis J (2007) Caffeinated beverage intake and the risk of heart disease mortality in the elderly: a prospective analysis. *Am J Clin Nutr* 85:392–398
- Pimentel GD, Zemdegs JC, Theodoro JA, Mota JF (2009) Does long-term coffee intake reduce type 2 diabetes mellitus risk? *Diabetol Metab Syndr* 1:6

6. Oba S, Nagata C, Nakamura K, Fujii K, Kawachi T, Takatsuka N, Shimizu H (2010) Consumption of coffee, green tea, oolong tea, black tea, chocolate snacks and the caffeine content in relation to risk of diabetes in Japanese men and women. *Br J Nutr* 103:453–459
7. Huxley R, Lee CM, Barzi F, Timmermeister L, Czernichow S, Perkovic V, Grobbee DE, Batty D, Woodward M (2009) Coffee, decaffeinated coffee, and tea consumption in relation to incident type 2 diabetes mellitus: a systematic review with meta-analysis. *Arch Intern Med* 169:2053–2063
8. van Dam RM, Hu FB (2005) Coffee consumption and risk of type 2 diabetes: a systematic review. *JAMA* 294:97–104
9. Matsuzawa Y (2005) Adiponectin: identification, physiology and clinical relevance in metabolic and vascular disease. *Atheroscler Suppl* 6:7–14
10. Weyer C, Funahashi T, Tanaka S, Hotta K, Matsuzawa Y, Pratley RE, Tataranni PA (2001) Hypoadiponectinemia in obesity and type 2 diabetes: close association with insulin resistance and hyperinsulinemia. *J Clin Endocrinol Metab* 86:1930–1935
11. Williams CJ, Fargnoli JL, Hwang JJ, van Dam RM, Blackburn GL, Hu FB, Mantzoros CS (2008) Coffee consumption is associated with higher plasma adiponectin concentrations in women with or without type 2 diabetes: a prospective cohort study. *Diabetes Care* 31:504–507
12. Kempf K, Herder C, Erlund I, Kolb H, Martin S, Carstensen M, Koenig W, Sundvall J, Bidel S, Kuha S, Tuomilehto J (2010) Effects of coffee consumption on subclinical inflammation and other risk factors for type 2 diabetes: a clinical trial. *Am J Clin Nutr* 91:950–957
13. Ruhl CE, Everhart JE (2005) Coffee and caffeine consumption reduce the risk of elevated serum alanine aminotransferase activity in the United States. *Gastroenterology* 128:24–32
14. Klatzky AL, Morton C, Udaltsova N, Friedman GD (2006) Coffee, cirrhosis, and transaminase enzymes. *Arch Intern Med* 166:1190–1195
15. Balk L, Hoekstra T, Twisk J (2009) Relationship between long-term coffee consumption and components of the metabolic syndrome: the Amsterdam Growth and Health Longitudinal Study. *Eur J Epidemiol* 24:203–209
16. Richard D, Kefi K, Barbe U, Poli A, Bausero P, Visioli F (2009) Weight and plasma lipid control by decaffeinated green tea. *Pharmacol Res* 59:351–354
17. Potenza MA, Marasciulo FL, Tarquinio M, Tiravanti E, Colantuono G, Federici A, Kim JA, Quon MJ, Montagnani M (2007) EGCG, a green tea polyphenol, improves endothelial function and insulin sensitivity, reduces blood pressure, and protects against myocardial I/R injury in SHR. *Am J Physiol Endocrinol Metab* 292:E1378–E1387
18. Hsu CH, Tsai TH, Kao YH, Hwang KC, Tseng TY, Chou P (2008) Effect of green tea extract on obese women: a randomized, double-blind, placebo-controlled clinical trial. *Clin Nutr* 27:363–370
19. Nagao T, Meguro S, Hase T, Otsuka K, Komikado M, Tokimitsu I, Yamamoto T, Yamamoto K (2009) A catechin-rich beverage improves obesity and blood glucose control in patients with type 2 diabetes. *Obesity* 17:310–317
20. Ryu OH, Lee J, Lee KW, Kim HY, Seo JA, Kim SG, Kim NH, Baik SH, Choi DS, Choi KM (2006) Effects of green tea consumption on inflammation, insulin resistance and pulse wave velocity in type 2 diabetes patients. *Diabetes Res Clin Pract* 71:356–358
21. van Dieren S, Uiterwaal CSPM, van der Schouw YT, van der A DL, Boer JMA, Spijkerman A, Grobbee DE, Beulens JWJ (2009) Coffee and tea consumption and risk of type 2 diabetes. *Diabetologia* 52:2561–2569
22. Odegaard AO, Pereira MA, Koh WP, Arakawa K, Lee HP, Yu MC (2008) Coffee, tea, and incident type 2 diabetes: the Singapore Chinese Health Study. *Am J Clin Nutr* 88:979–985
23. Kato M, Noda M, Inoue M, Kadowaki T, Tsugane S (2009) Psychological factors, coffee and risk of diabetes mellitus among middle-aged Japanese: a population-based prospective study in the JPHC study cohort. *Endocr J* 56:459–468
24. Kotani K, Tsuzaki K, Sano Y, Maekawa M, Fujiwara S, Hamada T, Sakane N (2008) The relationship between usual coffee consumption and serum C-reactive protein level in a Japanese female population. *Clin Chem Lab Med* 46:1434–1437
25. Arsenault BJ, Earnest CP, Despres JP, Blair SN, Church TS (2009) Obesity, coffee consumption and CRP levels in postmenopausal overweight/obese women: importance of hormone replacement therapy use. *Eur J Clin Nutr* 63:1419–1424
26. van Dam RM (2008) Coffee consumption and risk of type 2 diabetes, cardiovascular diseases, and cancer. *Appl Physiol Nutr Metab* 33:1269–1283
27. Hamer M, Witte DR, Mosdol A, Marmot MG, Brunner EJ (2008) Prospective study of coffee and tea consumption in relation to risk of type 2 diabetes mellitus among men and women: the Whitehall II study. *Br J Nutr* 100:1046–1053
28. Kagami K, Morita H, Onda K, Hirano T, Oka K (2008) Protective effect of caffeine on streptozotocin-induced beta-cell damage in rats. *J Pharm Pharmacol* 60:1161–1165
29. van Dijk AE, Olthof MR, Meeuse JC, Seebus E, Heine RJ, van Dam RM (2009) Acute effects of decaffeinated coffee and the major coffee components chlorogenic acid and trigonelline on glucose tolerance. *Diabetes Care* 32:1023–1025
30. Greenberg JA, Axen KV, Schnoll R, Boozer CN (2005) Coffee, tea and diabetes: the role of weight loss and caffeine. *Int J Obes* 29:1121–1129
31. van Dam RM, Feskens EJ (2002) Coffee consumption and risk of type 2 diabetes mellitus. *Lancet* 360:1477–1478